

TestHorse

Certified IT practice exam authority

Accurate study guides, High passing rate!
Testhorse provides update free of charge in one year!



Exam : AACD

**Title : American Academy of
Cosmetic Dentistry**

Version : DEMO

1. Why do self etch bonding systems tend to have less post operative sensitivity?

- A. Self etch bonding does not remove the smear layer but modifies it, leaving the smear plugs intact
- B. Self etch products remove & replace the smear layer
- C. Self etch bonding removes collagen fibrils and replaces them with resin plugs
- D. Self etch systems etch and bond simultaneously

Answer: A

2. In restoring multiple anterior teeth with ceramic restorations, the desired incisal plane can BEST be communicated to the laboratory by:

- A. Paralleling the incisal edges of the preparations.
- B. Indicating the incisal plane on the diagnostic cast.
- C. Including the patient's photograph with the lab prescription.
- D. Paralleling the unprepared mandibular incisal plane.

Answer: B

3. There is a direct relationship between the f-stop and depth of field.

Which is correct?

- A. Larger f-stop, larger depth of field
- B. Larger f-stop, smaller depth of field
- C. F-stop equals depth of field
- D. F-stop decreases depth of field

Answer: A

4. What is the BEST property of condensable composites?

- A. Lower filler content to allow lateral and vertical flow ensuring intimate contact
- B. Less stickiness and stiffer consistency and have a wear ratio similar to amalgam
- C. A filler loading of fibrous, porous, or irregular particles, and a filler percentage of 40% to 50% filled
- D. Clinical evidence of strength and polymerization shrinkage superior to conventional composites

Answer: D

5. A patient would like to brighten her smile in addition to restoring her missing tooth #8. However, she is not interested in dental implants. She would also like to close the spaces between her existing teeth.

Unfortunately, due to financial concerns she can only restore teeth ##7-10.

Reviewing the photograph above, how would you achieve golden proportion?

- A. Fabricating a three-unit bridge and a single veneer on tooth #10 will easily achieve golden proportion
- B. In office bleaching along with direct bonding on teeth ##7, 9, and 10, and a Maryland bridge to replace tooth #8 will easily achieve golden proportion
- C. Direct resin bonding in conjunction with a removable partial denture will achieve golden proportion
- D. It is impossible to achieve golden proportion unless more teeth are restored

Answer: A